AFOSR - TR - 76 - 1109

Approved for public release distribution unlimited.

COLORADO STATE UNIVERSITY

Fort Collins, Colorado 80523

July 31, 1976

TO: AIR FORCE OFFICE OF SCIENTIFIC RESEARCH

FINAL REPORT

ON

ELECTROCHEMICAL STUDIES IN

ALUMINUM CHLORIDE MELTS

AFOSR-75-2776

March 1, 1975 - May 31, 1976

OCT 19 1978

Principal Investigator: Dr. R. A. Osteryoung Department of Chemistry Colorado State University Fort Collins, Colorado 80523

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical real blanch reviewed and is
approved law paid on land TAN AFR 190-12 (7b).
Distribution is undensited.
A. D. BLAM
Technical information Officer

Market and the second second second

ACCESSION IN		
NTIS	White Sestion	
996	Butt Section	
UNANNOUNCED		£3
JUSTIFICATION		
,	*****************************	
F.A.		
	LANGILARGEITY CO	DE\$
Bisi.	es é leic	la:

### ELECTROCHEMICAL STUDIES IN ALUMINUM CHLORIDE MELTS

#### **ABSTRACT**

A

Work carried out under AFOSR-71-1995 was continued. The acid-dependent chemistry of sulfur was investigated. The Nernst behavior of the S/S couple was in accord with

$$S_2 + 4e = 2S^{=}$$

and the acid-base dependence can be expressed as

This explains the solubilization of metal sulfides in acid-chloroaluminate systems.

Oxides, selenides and tellurides show much the same behavior as sulfides. The stoichiometry indicates that the neutralization reaction for these bases may be written

where X is S, O, Se or Te.

A detailed electrochemical study of selenium has been undertaken. The following mechanism can be proposed:

Se + 3C1 
$$\stackrel{?}{=}$$
 SeC1  $\stackrel{?}{_3}$  + 2e  
SeC1  $\stackrel{?}{_3}$  + 2C1  $\stackrel{?}{=}$  SeC1  $\stackrel{?}{_5}$  + 2e

Selenium also appears to exist, as determined potentiometrically, as a dimer.

The electrochemistry of various molybdenum species has been studied. Mo(V) reacts to form Mo(III) which is further reduced to Mo(II) and Mo(0). The last two steps involve considerable surface chemistry. An E.C.E. mechanism is postulated to explain the Mo(V) +  $2e \Rightarrow Mo(III)$  electrode process.

OCT 19 1976

The work reported here is a 15-month continuation of activity reported previously under the Final Report to AFOSR-71-1995. Frequent reference will therefore be made to that Report.(1)

The work on sulfur and sulfides reported in (1) was continued, and it was demonstrated that the  ${\rm S/S}^{=}$  potential equilibrium could best be described by the reaction

$$S_2 + 4e \stackrel{?}{\leftarrow} 2S^{**}$$
 (1)

This was demonstrated by measuring the potential of an inert electrode in the AlCl<sub>3</sub>: NaCl melt at 175°C and the S/S<sup>=</sup> ratio was changed by coulometric oxidation of S<sup>=</sup> (for reduction of S). A Nernst plot gave linear behavior for a plot of E vs  $\log \frac{[S_2]}{[S^=]^2}$  assuming

$$[S_2] = [S]_T/2$$

The slope was  $\frac{2.3RT}{4F}$ , as per equation (1).

following this, the potential of the  $S_2/S^-$  couple was measured as a function of melt acidity. The results indicated that the previously reported (1) solubility of metal sulfides in the acid melt could best be explained by an equilibrium

$$A1SC1 + 3C1^{-} \neq S^{-} + A1C1_{\Delta}^{-}$$
 (2)

In the acid melt this equilibrium is displaced markedly to the left, markedly lowering S<sup>±</sup> activity. The results of this work were prepared for publication and are currently in press in the Journal of the American Chemical Society.

Further investigations have shown, as might be expected, that the solubilization shown by the metal sulfides is also observed for metal oxides, selenides and tellurides. These solubilities were determined from the height of the Cu<sup>+</sup> reduction cyclic voltammetry peak after equilibrating solid cuprous oxide, selenide or telluride with a 60:40 (AlCl<sub>3</sub>:NaCl) mole % melt. The values were all found to be similar at  $3.3 \pm 0.5 \times 10^{-1}$  mole liters<sup>-1</sup>, in agreement with the value determined earlier for cuprous sulfide. (2) It was suggested in

our previous report that this solubilization is brought about by some form of acid-base reaction between the melt and the chalcogenide ion. The potentiometric studies described above confirmed the nature of that reaction.

In order to investigate further the possible acid-base chemistry, it was decided to see if the chalcogenide ions could be titrated with  $\mathrm{Al}_2\mathrm{Cl}_7^-$  (the pricinpal acid species in the melt).(3) A 50:50 melt was prepared by anodizing an Al wire in an NaCl saturated melt until the required composition was reached. A small sample of  $\mathrm{Na}_2\mathrm{S}$ ,  $\mathrm{Na}_2\mathrm{Se}$  or  $\mathrm{Na}_2\mathrm{Te}$  was then added to the melt and the potential of an Al indicator electrode in the melt was observed to fall. The melt was then titrated with  $\mathrm{Al}^{3+}$  (from the anodization of an Al wire) using an Al indicator electrode. We also wished to observe the behavior of the oxide ion but, as has been observed previously,(4),  $\mathrm{Na}_2\mathrm{O}$  is not soluble in the melt. Instead we found that oxide ions could be introduced by adding  $\mathrm{BaCO}_3$  to the melt. Initially carbon dioxide was evolved and the residue,  $\mathrm{BaO}$ , was found to be soluble. The titration curves all showed the behavior typical of a strong acid-strong base reaction and the stoichiometry showed one chalcogenide ion to be equivalent to three  $\mathrm{Al}_2\mathrm{Cl}_7^-$  ions, indicating a reaction

$$x^{*} + 3A1_{2}C1_{7}^{-} \stackrel{K}{\neq} A1XC1 + 5A1C1_{4}^{-}$$
 (3)

where X is 0, S, Se or Te. All the species AlOC1, AlSC1, AlSeC1 and AlTeC1 have been prepared (5,6,7,8) by reacting metal chalcogenides with AlC1<sub>3</sub> in sealed tubes at elevated temperatures. In the future it is hoped to determine if these, or any other, species can be extracted from the melt. It is also hoped to be able to determine the magnitude of the equilibrium constant K. An order-of-magnitude calculation for the equilibrium constant for Equation 2 has been made. (10)

Apart from one very cursory study (9) (one cyclic voltammogram), the electrochemical behavior of selenium in these melts does not appear to have

been investigated. The lower oxidation states of selenium have, however, been the subject of a detailed spectroscopic study (10) and a number of polynuclear species with formal oxidation state less than unity have been identified in the eutectic melt as stable. We have recently commenced a detailed electrochemical study of selenium and various selenium species. In basic melts, pCl less than 3.5 (mole fraction scale), selenium shows a two-electron quasi-reversible reduction to selenide and two quasi-reversible two-electron oxidations (the number of electrons being determined coulometrically by exhaustive controlled potential electrolysis). From the pCl dependence of the cyclic voltammetry peak potentials for the oxidations, the following mechanism for the oxidation of selenium in basic melts can be proposed

Se + 3C1 
$$\stackrel{?}{=}$$
 SeC1 $\stackrel{?}{_3}$  + 2e  $\stackrel{?}{=}$  (4)

$$SeCl_3^- + 2Cl^- \neq SeCl_5^- + 2e^-$$
 (5)

When the pCl of the melt is increased to greater than about 3.5 selenium still shows a two-electron reduction but the oxidation now becomes a single four-electron process, again quasi-reversible. The potential of this oxidation now shows no acidity dependence, implying that the  $Se^{4+}$  species is not solvated by any acidic or basic species (Cl<sup>-</sup>, Al<sub>2</sub>Cl<sub>7</sub> or AlCl<sub>3</sub>). Support for these proposed oxidation mechanisms is at present being sought from spectroscopic studies.

Equilibrium potential measurements of the selenium-selenide couple have been made in both acidic and basic melts and also for fused selenium and selenide concentration as a function of acidity. The measurements for variable selenium and selenide concentrations indicate that in both acid and basic melts, selenium is present as a dimer, while a plot of equilibrium potential against pC1 for fixed selenium and selenide concentrations has a alope of  $\frac{3RT}{2F}$  (120 mV). From these observations the following electrode reaction for the reduction of selenium can be postulated

$$Se_2 + 2A1C1_4^- + 4e \neq 2A1SeC1 + 6C1^-$$
 (6)

$$A1SeC1 + 3C1^{-} \neq Se^{-} + A1C1_{A}^{-}$$
 (7)

with the latter equilibrium lying well to the left hand side. This is the same mechanism as was proposed for the reduction of sulfur (see above and 10) and is in agreement with our earlier postulates concerning the acid-base chemistry of metal chalcogenides in these melts. The kinetics of the selenium reduction also show marked acidity dependence and are being studied further.

The electrochemical work on the chemistry of molybdenum was continued utilizing the rotating ring-disc glassy carbon electrode.(1) The electrochemistry of Mo(V) and Mo(VI) was found to be identical; however, it was discovered that Mo(VI), introduced as  $MoO_3$ , oxidized the melt to yield  $Cl_2$ , being itself reduced to Mo(V). The Mo(V) reduces to Mo(III) in a diffusion controlled, first-order process. Further reduction, to the Mo(II) and Mo(0) state is very complex and involves considerable surface chemistry. Mo(V) could be quantitatively reduced to Mo(III); however, the ring-disc collection efficiency for this process was rotation rate dependent, increasing with decreasing rotation rate. This indicates that a chemical step, perhaps the generation of an initially electrochemically inactive species, of Mo(V), occurs or an ECE mechanism exists.

A possible scheme to explain the Mo(III)-Mo(V) electrochemistry is

$$Mo(III) \stackrel{?}{=} Mo(IV) + e \qquad E$$
 $Mo(IV) \stackrel{?}{=} Mo'(IV) \qquad C \qquad \qquad (8)$ 
 $Mo'(IV) \stackrel{?}{=} Mo(V) + e \qquad E$ 

The cyclic voltammetric and ring-disc work tends to support this scheme, since values of  $\frac{1}{\sqrt{1/2}}$  and  $\frac{1}{\sqrt{1/2}}$  are close to 2:1 for low:high sweep rates or rotation rates.

The Mo(III)-Mo(V) chemistry exhibited marked acidity dependence. The overall scheme, involving the measurement of the Mo(III)-Mo(V) potential as the melt acidity was varied, is consistent with

$$Mo(III) + 2C1^{-} \neq Mo(V) + 2e$$
 (9)

The detailed mechanism, however, is complex and is incomplete at this time. Clearly,  $Cl^-$  must be involved in the E.C.E. scheme. The chemical step, given above as  $Mo(IV) \neq Mo^+(IV)$ , may be a second order disproportionation - i.e.,

$$2Mo(IV) \stackrel{?}{\leftarrow} Mo(III) + Mo(V) \tag{10}$$

Some experimental data appears to support this conclusion.

The very complex nature of Mo-chemistry is clear. Earlier work indicating that  ${\rm MoO_3}$  yields either  ${\rm MoO_2Cl_2}$  appears incorrect(11), although the oxidation of the melt by  ${\rm MoO_3}$  is slow and may be acid dependent.

It is hoped that the problem of Mo-electrochemistry may be completed in the near future.

During the period of this report several meetings were held with personnel at the Seiler Research Laboratory at the United States Air Force Academy. Discussions were held relating to both the sodium tetrachloroaluminate system and the room temperature ethylpyridinium bromide-aluminum chloride melt. In addition, Lieutenant Charles Hussey spent a week or so working in the laboratory at Colorado State. Numerous telephone contacts were maintained with both Hussey and Colonel Lowell King of the Seiler Laboratory.

### REFERENCES

- 1. R. A. Osteryoung, Final Report, AFOSR-17-1995, Colorado State University, Fort Collins, CO 80523, June 30, 1975.
- 2. L. G. Boxall, H. L. Jones and R. A. Osteryoung, J. Electrochem. Soc., 120, 223 (1973).
- 3. G. Letisse and B. Tremillon, J. Electroanal. Chem., 17, 387 (1968).
- 4. H. Kuhnl and U. Geffarth, Zanorg. allg. Chem., 391, 280 (1972).
- 5. P. Hagenmuller, J. Rouxel, J. David, A. Colin and B. LaNeidr, Z. Anorg. Allg. Chem., 323, 1 (1963).
- 6. J. Rouxel and P. Palvadeau, Bull. Soc. Chim. France, 2698 (1967).
- 7. J. Rouxel and P. Palvadeau, Bull. Soc. Chim. France, 2044 (1966).
- 8. R. Marassi, G. Mamantov and J. Q. Chamber, Inorg. Nucl. Chem. Letts., 11, 245 (1975).
- 9. R. Fehrmann, N. J. Bjerrum and M. A. Andreasen, Inorg. Chem., <u>14</u>, 2259 (1975).
- 10. K. A. Paulsen and R. A. Osteryoung, J. Amer. Chem. Soc., in press.
- 11. H. Kuhni, Angew. Chem., Int. Ed., <u>6</u>, 641 (1967).

Publications -- Grant-related Activity (includes activity referred to in Reference 1)

- Janet Osteryoung and R. A. Osteryoung, "The Advantage of Charge Measurements for Determining Kinetic Parameters", Electrochimica Acta 16, 525 (1971).
- R. A. Osteryoung, "Introduction to the On-Line Use of Computers in Electro-chemistry", Vol. II, "Application of Computers to Chemical Instrumentation", Ed. by Mattson, Mark and MacDonald, MARCEL DEKKER (1973).
- L. G. Boxall, H. L. Jones and R. A. Osteryoung, "Solvent Equilibria in AlCl NaCl Melts", J. Electrochem. Soc.,  $\underline{120}$  (2), 223 (1973).
- H. Lloyd Jones, L. G. Boxall and R. A. Osteryoung, "Organic Electrode Reactions in Fused AlCl $_3$  Containing Solvents", J. Electroanal. Chem.,  $\underline{38}$ , 476 (1972).
- L. G. Boxall, H. L. Jones and R. A. Osteryoung, "Electrochemical Studies on Ag, Fe and Cu Species in AlCl $_3$ -NaCl Melts", J. Electrochem. Soc.,  $\underline{121}$ , 212 (1974).
- H. Lloyd Jones and R. A. Osteryoung, "Electrode Reactions of Aromatic Amines in Solvents Containing Fused AlCl<sub>3</sub>, II.", J. Electroanal. Chem.,  $\underline{49}$ , 281 (1974).
- R. J. Gale and R. A. Osteryoung, "Investigation of Subvalent Ion Effects During Aluminum Anodization in Molten NaCl-AlCl<sub>3</sub> Solvents", J. Electrochem. Soc., 121 983 (1974).
- V. R. Koch, L. L. Miller and R. A. Osteryoung, "Reductive Defunctionalization of 1-substituted Adamantanes in Molten Sodium Tetrachloroaluminate", J. Org. Chem.,  $\underline{39}$ , 2416 (1974).
- H. Lloyd Jones and R. A. Osteryoung, "Organic Reactions in Molten Tetrachloroaluminate Solvents", Advances in Molten Salt Chemistry, Vol. 3, Edited by J. Braunstein, G. P. Smith and G. Mamantov, Plenum Publishing (1975).
- R. J. Gale and R. A. Osteryoung, "Dissociative Chlorination of Nitrogen Oxides and Oxyanions in Molten Sodium Chloride-Aluminum Chloride Solvents", Inorg. Chem., 14, 1232 (1975).
- H. L. Chum, V. R. Koch, L. L. Miller and R. A. Osteryoung, "An Electrochemical Scrutiny of Organometallic Iron Complexes and Hexamethylbenzene in a Room Temperature Molten Salt", J. Am. Chem. Soc., <u>97</u>, 3264 (1975).
- D. E. Bartak and R. A. Osteryoung, "The Electrochemical Oxidation of N,N,N',N'-Tetramethylbenzidine in Molten Sodium Tetrachloroaluminate", J. Electrochem. Soc., 122, 600 (1975).
- J. Phillips, R. J. Gale, R. G. Wier and R. A. Osteryoung, "Glassy Carbon Rotating Ring-Disc Electrodes for Molten Salt Studies", Anal. Chem., 48, 1266 (1976).
- D. E. Bartak and R. A. Osteryoung, "The Redox Behavior of Tetrachloro-p-Benzoquinone Tetrachlorohydroquinone Systems in Molten Aluminum Chloride-Sodium Chloride Solvents", J. Electroanal. Chem., in press (1976).
- V. R. Koch, L. L. Miller and R. A. Osteryoung, "Electroinitiated Friedel-Crafts Transalkylations in a Room Temperature Molten Salt Medium", J. Am. Chem. Soc., in press (1976).

- K. A. Paulsen and R. A. Osteryoung, "Electrochemical Studies on Sulfur and Sulfides in AlCl<sub>3</sub>-NaCl Melts", J. Am. Chem. Soc., in press (1976).
- R. A. Osteryoung, "Chemistry and Electrochemistry in Aluminum Chloride Molten Salt Systems", Symposium on Molten Salts, Symposium Volume, The Electrochemical Society, in press (1976).

# Manuscripts in Preparation -- Related to Grant Activity

- J. Phillips and R. A. Osteryoung, "Electrochemical Studies of Mo-Species in AlCl<sub>3</sub>-NaCl Solvent Systems", to be submitted to J. Am. Chem. Soc., or J. Electrochem. Soc.
- J. Robinson and R. A. Osteryoung, "Acid-Base Chemistry of Chalcogenides in Sodium Tetrachloroaluminate Melts", to be prepared for J. Am. Chem. Soc.

# Meeting Presentation and Seminars--Related to Grant Activity

- R. A. Osteryoung, "Computerized Electrochemical Experimentation", Dept. of Chemistry Colloquia, Pennsylvania State University, University Park, PA, May 27, 1971.
- L. G. Boxall, H. L. Jones and R. A. Osteryoung, "Electrochemical Studies in Aluminum Chloride Melts", N.W. Regional ACS Meeting, Bozeman, Montana, June, 1971.
- R. A. Osteryoung, "Application of a Computer-Based Pulse Polarographic System in Molten Salt Studies", Gordon Research Conference on Molten Salts, Kimball Union Academy, Meriden, N.H., August, 1971.
- R. A. Osteryoung, "Use of a Mini-Computer in Electrochemical Studies", Symposium on Mini-Computers in the Research & Teaching Laboratory, American Chemical Society Meeting, Washington, D.C., September, 1971.
- R. A. Osteryoung, "Chemistry and Electrochemistry in Fused Salts", Baylor University (ACS Waco Station), May 8, 1972.
- R. A. Osteryoung, "Computer Controlled Electrochemical Experimentation", Texas Christian University (Dallas-Ft. Worth ACS Section), May 9, 1972.
- R. A. Osteryoung, "Computer Controlled Electrochemical Experimentation", (San Antonio, Texas ACS Section), May 10, 1972.
- R. A Osteryoung, "Computer Controlled Electrochemical Experimentation", Texas A a H (College Station, Texas ACS Section), May 11, 1972.

Electric Control

- R. A. Osteryoung, "Computer Controlled Electrochemical Experimentation", Midwestern University (Wichita Falls, Texas/Duncan, Okla. ACS Section), May 12, 1972.
- H. L. Jones, L. G. Boxall and R. A. Osteryoung, "Organic Electrochemistry in Aluminum Halide Melts", Rocky Mountain Regional ACS Meeting, Ft. Collins, CO, June 1972.
- R. A. Osteryoung, H. L. Jones and L. G. Boxall, "Electrochemical Studies in Molten Chloroaluminates", Symposium on Fused Salt Technology, Electrochemical Society Meeting, Chicago, May 8-13, 1973.
- R. A. Osteryoung, R. H. Abel, L. G. Boxall and B. H. Vassos, "An Introduction to the On-Line Use of Digital Computers in Electrochemistry", Plenary Lecture at Symposium on Electrochemical Measurements, by Digital Computer, Electrochemical Society Meeting, Chicago, May 8-13, 1973.
- D. E. Bartak and R. A. Osteryoung, "Oxidation of Tetramethylbenzidine in Chloroaluminate Melts", Electrochemical Society, San Francisco, CA, May, 1974.
- R. A. Osteryoung, "Chemistry in Aluminum Chloride Melts", Fifth International Conference in Non-Aqueous Solutions, International Union of Pure and Applied Chemistry, Vienna, July 10-12, 1974.
- R. A. Osteryoung, "Electrochemical Studies in Fused Salts", First Latin American Electrochemistry and Corrosion Meeting (ABRACO), Rio de Janeiro, Brazil, Oct. 21-25, 1974.
- R. A. Osteryoung, "Some Applications of Pulse Techniques to Analytical Chemistry and Electrochemistry", University of Brussels, Brussels, Belgium, March 17, 1975.
- V. R. Koch, L. L. Miller and R. A. Osteryoung, "Electroinitiated Friedel-Crafts Transalkylation in a Room Temperature Molten Salt Medium", Mile High Electrochemistry Symposium, Colorado Electrochemical Society Section, Fort Collins, CO, May 3, 1975.
- K. A. Paulsen and R. A. Osteryoung, "Electrochemical Studies of Sulfur in Molten Sodium Tetrachloroaluminate", Fall Meeting, American Chemical Society, Chicago, Illinois, August 1975.
- R. A. Osteryoung, "Chemistry and Electrochemistry in Aluminum Chloride Molten Salt Systems", International Symposium on Molten Salts, Spring Meeting, The Electrochemical Society, Washington, D. C., May 2-7, 1976.
- R. A. Osteryoung, "Electrochemical Studies in Molten Sodium Chloride-Aluminum Chloride", Colorado College, Colorado Springs, Colorado, December 4, 1975.
- R. A. Osteryoung, "Electrochemical Studies in Molten Chloroaluminates", Gould, Inc., Rolling Meadows, Illinois, March 4, 1976.
- R. A. Osteryoung, "Chemical and Electrochemical Studies in Molten Chloro-aluminate Solvents", Dept. of Chemistry Colloquium, Purdue University, W. Lafayette, Indiana, March 23, 1976.
- V. R. Koch, L. L. Miller and R. A. Osteryoung, "Electroinitiated Friedel-Crafts Transalkylations in a Room Temperature Molten Salt Medium", Symposium on Electroorganic Oxidations, Electrochemical Society Meeting, Washington, D.C., May 2-7, 1976.

AFOSRI - TR - 76 - 1109 FILL	READ INSTRUCTIONS
	BEFORE COMPLETING FORM
CITLE (and Subilitie)	1 M. C. C. SIA
	10 1 1 Mar / 311
XC.	TO THE OF REPORT OF PRINCIPLE
ELECTROCHEMICAL STUDIES IN ALUMINUM CHLORIDE	L
THE DISC.	FERFORMING ORG. REPORT NUMBER
7 AUTHOR(a)	S CONTRACT OR GRANT NUMBER(s)
Robert A. Osteryoung	AFOSR-75-2776 NEW
Marie and the second se	I NOW IS ZITE NO
9 PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Colorado State University	9536-01, 61102F, 681303
Fort Collins, Colorado 80523	
Air Force Office of Scientific Research/NC	ME HE-ONI DATE (12)
Building 410	Jule 1076
Bolling AFB, D.C. 20332  14 MONITORING AGENCY NAME & ADDRESS(II dillerent from Controlling Office)	10
	18. SECURITY CLASS, (of this
51). 111-11FOSK-2776-75 ( )	Unclassified
	184. DECLASSIFICATION/DOWNGRADING
16 DISTRIBUTION STATEMENT (of this Report)	1
Approved for public release; distribution unlimi	ted
TOTAL BELLIAME	141
1 00/ 11-9000 (11/900	3602 1
IS SUPPLEMENTARY NOTES	
19. KEY WORDS (Continue on reverse side if necessary and identity by block number) molten salts; sodium tetrachloroaluminate; sulfur	r; molybdenym;
chalcogenides	
20 ABSTRACT (Continue on reverse side if necessary and identify by block number) Work carried out under AFOSR-71-1995 was con	atinued. The acid dependent
chemistry of sulfur was investigated. The Nernst	
me nernst	r newayior of the 3/2 couple
was in accord with	
was in accord with	
S <sub>2</sub> + 4e = 25 <sup>=</sup>	
-	(OVED)
$S_2 + 4e = 2S^{\pm}$ and the acid-base dependence can be expressed as	(OVER)
S <sub>2</sub> + 4e = 2S <sup>±</sup> and the acid-base dependence can be expressed as  DD , FORM 1473 EDITION OF 1 NOV 6B IS OBSOLETE  S/N 0.102-0.44-0.001   Unc	lassified
S <sub>2</sub> + 4e = 25 <sup>±</sup> and the acid-base dependence can be expressed as  DD , FORM 1473 1473 EDITION OF 1 NOV 68 IS OBSOLETE  S/N 0.102-0.14-0.001   Unc	
S <sub>2</sub> + 4e = 25 <sup>±</sup> and the acid-base dependence can be expressed as  DD , FORM 1473 1473 EDITION OF 1 NOV 68 IS OBSOLETE  S/N 0.102-0.14-0.001   Unc	lassified

CUMITY CLASSIFICATION OF THIS PAGE(When Date Entered

S + A1C1 ≠ A1SC1 + 3C1

This explains the solubilization of metal sulfides in acid-chloroaluminate systems. The Phaneral

dxides, selenides and tellurides show much the same behavior as sulfides. The stoichiometry indicates that the neutralization reaction for these bases may be written

where X is S, O, Se or Te.

A detailed electrochemical study of selenium has been undertaken. The following mechanism can be proposed:

Se + 3C1 
$$\stackrel{?}{=}$$
 SeC1 $\frac{7}{3}$  + 2e  
SeC1 $\frac{7}{3}$  + 2C1  $\stackrel{?}{=}$  SeC1 $\frac{7}{5}$  + 2e

Selenium also appears to exist, as determined potentiometrically, as a dimer

The electrochemistry of various molybdenum species has been studied % Mo(V) reacts to form Mo(III) which is further reduced to Mo(II) and Mo(0). The last two steps involve considerable surface chemistry. The E.C.E. mechanism is postulated to explain the Mo(V) +  $2e \neq Mo(III)$  electrode process.